

CEREBRAL PALSY

V.K. Srivastava

N. Laisram

R.K. Srivastava

ABSTRACT

The clinical pattern and etiology of 544 cases of cerebral palsy were studied retrospectively. Of these cases, 354 (65.1%) were males. Four hundred and ninety seven (91.4%) cases were of spastic type. Hypotonic, ataxic and athetoid cerebral palsy were observed in 5.5, 1.5 and 1.3% cases, respectively. There was one case each of tremor and mixed type. In the spastic group, quadriplegia comprised the maximum number of cases (34.9%). Hemiplegia (28.7%) and diplegia (21.9) were also common. Mental retardation was found in 47.2%, while speech impairment was observed in 37% cases. Other handicaps included visual (9%), seizures (8.8%), and auditory handicap (2.9%). The etiological factors were prenatal in 7.7% cases, natal in 43.8% cases and postnatal in 26.1% cases. More than one etiological factor was observed in 14.5% cases, while in 7.9% cases, no apparent cause could be found.

Key words: Cerebral palsy, Etiology, Clinical profile.

From the Departments of Pediatrics and Physical Medicine and Rehabilitation, Safdarjang Hospital, New Delhi 110 029.

Reprint requests: Dr. V.K. Srivastava, Professor and Head of Unit, Department of Pediatrics, Safdarjang Hospital, New Delhi 110 029.

Received for publication: December 19, 1991;

Accepted: May 6, 1992

Cerebral palsy continues to be one of the major crippling disorders in children. Various studies(1-8) have been carried out to study the clinical types and the etiology of this disorder. However, the exact etiology of this condition is often not identified. The present study was done to find the clinical profile of cerebral palsy and identify the etiological factors.

Material and Methods

Five hundred and forty four cases of cerebral palsy attending the Pediatrics and Physical Medicine and Rehabilitation Departments of Safdarjang Hospital between 1981 and 1989 comprised the study group. The cases diagnosed as cerebral palsy included those having "persistent disorder of movement and posture appearing early in life and due to a developmental non progressive disorder of the brain"(9). The clinical types were classified as described by Mitchell(3), including the spastic, hypotonic, tremor, ataxic and mixed types. The spastic group was further subclassified as monoplegia, paraplegia, hemiplegia, quadriplegia, diplegia and triplegia. Etiological factors were grouped as prenatal, natal, postnatal, mixed and unknown. Prenatal factors included various causes resulting in brain damage from conception till onset of labor. Natal causes included various factors operating during birth process, while postnatal factors included those affecting the brain after birth. Where there was more than one factor, they were labelled as 'mixed'. Term babies with birth weight 2.5 kg or more and with normal prenatal, natal and postnatal history were labelled as of unknown origin. Convulsions were considered as causative factor in the postnatal group when the motor deficit followed the seizure episode. The associated handicaps were also noted.

Results

Of 544 cases, there were 354 males (65.1% cases) and 190 females (34.9% cases). The types of cerebral palsy are shown in *Table I*. Mental retardation of varying degrees was observed to be the most common handicap (257 cases). Mild, moderate and severe were observed in 149, 28 and 76 cases, respectively while four cases had profound degree of mental retardation. Visual defect, seizure and hearing impairment were present in 9, 8.8 and 2.9%, respectively. The visual defects present were strabismus, nystagmus, cataract and bilateral optic atrophy.

Maximum cases were due to natal causes (43.8%) followed by postnatal causes (26.1%). A total of 14.5% cases had more than one etiological factor, while in 7.9% cases there was no known cause. The various etiological factors in the prenatal, natal and postnatal groups are shown in *Table II*. In cases where more than one etiological factor was present, the most frequent causes were a combination of

prematurity or birth anoxia in association with antepartum hemorrhage, twin, forceps, or cesarean delivery.

Discussion

A male preponderance was observed with the male to female ratio being 1.86 : 1. This is consistent with other reports(4,6). Of the different types of cerebral palsy, spasticity was the most common. Our finding is in agreement with other workers(3,4,6,7) who have reported spasticity in 62.7 to 80% cases. Among the spastic type, quadriplegia was most common (34.9%) followed by hemiplegia (28.7%) and diplegia (21.9%). Basu(4) found a high proportion of cases with tetraplegia and diplegia while hemiplegia was found in a smaller number of cases. Mitchell(3) found hemiplegia to be most common (37.1%) followed by tetraplegia in 19.2% cases. O'Reilly and Walentynowicz(7) also found hemiplegia to be the most common (26.1%), followed by paraplegia (15.7%), while quadriplegia was found in only 13.5% cases.

In our study, hypotonia, ataxia, and athetosis were present in 5.6, 1.5 and 1.3% cases, respectively while tremor and mixed type was observed in one case each (0.2%). There was no case of rigidity in our study. O'Reilly and Walentynowicz(7) observed a higher incidence of mixed, athetoid, rigidity and ataxic cerebral palsy in 12.0, 11.7, 7.2 and 4.9% cases, respectively. The incidence of atonia and tremor was low, i.e., 1.1 and 0.3%, respectively. Mitchell(3) also observed higher incidence of mixed (9.6%) and athetoid (7.5%) types, while ataxia, hypotonia and tremor was observed in 1.7, 2.3 and 0.4% cases, respectively. Basu(4) observed hypotonia, athetosis and mixed in 7.1, 5.5 and 2.4% cases, respectively while no case of ataxia was found. Makwabe and

TABLE I—Types of Cerebral Palsy

Type	No (n = 544)	Percentage
Spasticity	497	91.4
Quadriplegia	190	34.9
Hemiplegia	156	28.7
Diplegia	119	21.9
Paraplegia	24	4.4
Monoplegia	5	0.9
Triplegia	3	0.6
Hypotonic	30	5.5
Ataxic	8	1.5
Athetoid	7	1.3
Tremor	1	0.2
Mixed	1	0.2

TABLE II—*Etiological Profile in Cerebral Palsy*

Prenatal (n = 42)	No (%)	Natal (n = 238)	No (%)	Postnatal (n = 142)	No (%)
Microcephaly	10 (1.8)	Anoxia	133 (24.5)	Encephalitis	65 (12)
Toxemia	7 (1.3)	Prematurity	22 (4.0)	Meningitis	28 (5.2)
Antepartum- hemorrhage	5 (0.9)	Prolonged labor	18 (3.3)	Convulsion	20 (3.7)
Hydrocephalus	3 (0.6)	Forceps	15 (2.8)	Head injury	13 (2.4)
Drugs	3 (0.6)	Breech	12 (2.2)	Acute infantile hemiplegia	8 (1.5)
Twins	3 (0.6)	Low birth weight	11 (2.0)	Neonatal jaundice	8 (1.5)
Diabetes	2 (0.4)	Postmaturity	9 (1.7)		
Consanguinity	1 (0.2)	Cesarean	8 (1.5)		
Rubella	1 (0.2)	Precipitate labor	5 (0.9)		
Toxoplasmosis	1 (0.2)	Cord around neck	3 (0.6)		
Other maternal infections	6 (1.1)	Face presentation	2 (0.4)		

Mgone(6) observed a higher incidence of athetosis (12%), while hypotonia and mixed type was seen in 4% cases each.

Mental retardation was the most common associated handicap (47.2%). Basu(4) also observed mental retardation to be the most common (71.6%) related disorder.

Out of 544 cases of cerebral palsy, majority of the cases (43.8%) were due to natal etiology. Our findings are in accordance with Perlstein(8) and O'Reilly and Walentynowicz(7) who found natal etiology to be the most common cause of cerebral palsy. The incidence of postnatal cause was high (72%) in the study of Makwabe and Mgone(6). In our study, postnatal causes were observed to be second most common (26.1%). Prenatal causes were least common (7.7%). O'Reilly and Walentynowicz(7) and Perlstein(8), however, found higher incidence of prenatal causes, i.e., 38.5 and 30%, respectively. The low

incidence of prenatal causes in our study may be due to lack of facilities for detection of factors in the prenatal period.

Of the prenatal factors, microcephaly (1.8%) and toxemia (1.3%) were the most common etiological factors. Among natal factors, anoxia was the most common (24.5% cases). Infections of the central nervous system comprised the major etiopathogenic factor of the postnatal causes. Encephalitis and meningitis were found in 12% and 5.2% cases, respectively. O'Reilly and Walentynowicz(7) in their study found multiple pregnancy and idiopathic factors to be the most frequent causes in the prenatal group, i.e., 5.6 and 1.9%, respectively. The most common natal etiology in their study was prematurity (22.7%), followed by anoxia (7.7%). In the postnatal group, encephalitis has been reported to be the most frequent factor (6.9%) which is similar to our observation.

The present study reveals that the natal and postnatal factors are responsible for more than two-thirds of cases of cerebral palsy. These results indicate the need to reinforce the existing maternal and child health services existing in the country. The study also highlights the multifaceted problems relating to cerebral palsy. The child with cerebral palsy may not be confined to the physical disability alone but may have other associated handicaps which must be detected at the earliest. This will facilitate a timely and appropriate intervention.

REFERENCES

1. Nelson KB, Ellenberg JH. Antecedents of cerebral palsy, Univariate analysis of risks. *Am J Dis Child* 1985, 139: 1031-1038.
2. Nelson KB, Ellenberg JH. Antecedents of cerebral palsy: A multivariate analysis of risks. *N Engl J Med* 1986, 315: 81-86.
3. Mitchell RG. The general medical and neurological features. *In: Cerebral Palsy in Childhood and Adolescence*. Ed Henderson JL. Edinburgh, E & S Livingstone Ltd, 1961, pp 37-72.
4. Basu B. Cerebral palsy in childhood. *J Indian Med Assoc* 1969, 49: 477-482.
5. Izuora GI, Okoro ABO. Some clinical aspects of cerebral palsy among Nigerian Ibo children. *Centr Afr J Med* 1981, 27: 155-159.
6. Makwabe CM, Mgone CS. The pattern and etiology of cerebral palsy as seen in Dar-es-Salam, Tanzania. *East Afr Med J* 1984, 12: 896-899.
7. O'Reilly D Elliot, Walentynowicz JE. Etiological factors in cerebral palsy: A historical review. *Dev Med Child Neur* 1981, 23: 633-642.
8. Perlstein MA. Infantile cerebral palsy—classification and clinical correlations. *JAMA* 1952, 149: 30-34.
9. Mackeith RC. The little club memorandum on terminology and classification of cerebral palsy. *Cerebral Palsy Bull* 1959, 27: 34-37.

NOTES AND NEWS

III MAHARASHTRA STATE CONFERENCE OF IAP

The III Maharashtra State Conference of IAP is to be held on *21st and 22nd November, 1992* at Nasik.

Registration Fees upto 30th September, 1992 are Rs. 250/- and Rs. 300/- thereafter, to be paid by demand draft payable at Nasik.

For further details, please contact:

Dr. S.S. Kashyape,
Nagjee Memorial Hospital,
Wadala Road,
Nasik 422 001.
Tel: 73201/74780