

**WEB TABLE I** UNIVARIATE ANALYSIS OF PUTATIVE RISK FACTORS FOR MORTALITY

<i>Risk factors</i>	<i>Died</i> (n=71)	<i>Survived</i> (n=78)	<i>Odds ratio</i> (95% CI)	<i>P value</i>
Gestational age ≥28 wks	38	17	4.13 (2.03-8.42)	<0.001
Birth weight ≥800g	35	22	2.48 (1.26-4.88)	0.008
Male	37	41	0.98 (0.52-1.87)	0.96
SGA	28	49	0.39 (0.19-0.75)	0.004
Adequately supervised antenatally	12	15	1.17 (0.51-2.71)	0.71
Abnormal antenatal doppler present	13	24	0.50 (0.23-1.09)	0.079
Received antenatal steroids	49	62	0.58 (0.27-1.21)	0.14
Asphyxia present	20	10	2.67 (1.15-6.19)	0.02
Age of transfer ≥1 hr	60	59	1.76 (0.77-4.01)	0.18
Received surfactant	38	18	3.84 (1.89-7.76)	<0.001
Any ventilation	65	34	14.02 (5.43-36.19)	<0.001
Invasive ventilation	53	16	11.41 (5.3-24.56)	<0.001
Pneumothorax	2	1	0.45 (0.04-5.05)	0.51
PPHN	6	0	2.2 (1.84-2.63)	0.009
Shock	52	11	16.67 (7.29-38.08)	<0.001
Any sepsis <sup>+</sup>	66	61	3.68 (1.28-10.58)	0.01
Culture positive sepsis	21	29	0.71 (0.36-1.41)	0.33
NEC	20	14	1.79 (0.83-3.89)	0.14
Anemia requiring blood transfusion	40	39	1.29 (0.68-2.46)	0.44
Presence of PDA	30	21	1.99 (0.99-3.95)	0.049

Values are n. <sup>+</sup>suspected and culture positive sepsis. PPHN – persistent pulmonary hypertension of newborn, SGA- small for gestational age, PDA-hemodynamically significant patent ductus arteriosus. p<0.05 significant.