

WEB TABLE I BIVARIATE ANALYSIS OF CLINICAL FEATURES AND DETERMINATION OF ETIOLOGY

<i>Clinical feature</i>	<i>Diagnosed N(%)</i>	<i>Undiagnosed n(%)</i>	<i>Odds ratio</i>	<i>95% CI</i>	<i>P value</i>	<i>+LR</i>	<i>-LR</i>
Regression of milestones	6 (7.2)	6 (33.3)	0.1558	0.0431-0.5632	0.007	0.21	1.39
Behavioural problem	9 (10.8)	6 (33.3)	0.2432	0.0733-0.8073	0.025	0.32	1.337
Seizures	29 (34.9)	10 (55.6)	0.4296	0.1529-1.2076	0.058	0.62	1.466
Consanguinity	10 (12)	4 (22.2)	0.4795	0.1316-1.7468	0.217	0.54	1.13
Family history	15 (18.1)	1 (5.6)	3.75	0.4625-30.4061	0.17	3.23	0.86
Microcephaly	26 (31.3)	4 (22.2)	1.5965	0.4789-5.3225	0.23	1.4	0.88
Facial dysmorphism	37 (44.6)	8 (44.4)	1.0054	0.3606-2.8037	0.49	1.004	0.99
Non facial dysmorphism	36 (43.4)	5 (27.8)	1.9915	0.6504-6.0978	0.117	1.56	0.78
Neurological deficit	42 (50.6)	6 (33.3)	2.0488	0.7025-5.9748	0.098	1.51	0.74

LR: Likelihood ratio, CI: Confidence interval.