

WEBTABLE I GEOMETRIC MEAN CONCENTRATIONS (GMC) OF ROTAVIRUS-SPECIFIC FECAL, SALIVARY AND SERUM IGA IN INFANTS GIVEN THREE OR FIVE DOSES OF ROTAVIRUS VACCINES

<i>Serum IgA response categorization</i>	<i>Rotavirus-specific fecal IgA U</i>						<i>Rotavirus-specific serum IgA</i>			<i>Rotavirus-specific salivary IgA</i>		
	<i>3 dpv Dose 1 (n=42)</i>	<i>28 dpv Dose1 (n=42)</i>	<i>3 dpv Dose2 (n=25)</i>	<i>28 dpv Dose2/4 (n=25)</i>	<i>3-7 dpv Dose3/5 (n=41)</i>	<i>pre-post fold change</i>	<i>Pre vaccination (n=42)</i>	<i>28 dpv (n=42)</i>	<i>pre-post fold change</i>	<i>Pre vaccination (n=42)</i>	<i>28 dpv (n=42)</i>	<i>pre-post fold change</i>
No response (n=13)	38.7	27.8	24.7	20.9	18.3	0.5	1	2	1.5	7.5	1.6	0.2
<2 fold (n=4)	74.1	71.6	ND*	ND*	315.9	4.3	56.4	93.9	1.7	27.5	1.8	0.1
2-4 fold (n=6)	35.1	16.4	11.5	15.5	40.7	1.2	22.6	63.9	2.8	7.7	9.6	1.3
≥4 fold (n=19)	56.8	56.0	17.6	34.5	116.3	2.0	19.9	265.5	13.4	1.4	8.2	5.6

*ND: Fecal IgA measurement was not done in samples from this group; dpv: days post vaccination.