

WEB TABLE II PREGNANCY-RELATED RISK FACTORS OF INTRA-PARTUM RELATED PERINATAL MORTALITY (IPPM)

<i>Variable</i>	<i>IPPM + (n=20)</i>	<i>IPPM – (n=228)</i>	<i>P value</i>
Parity	1 (1-2)	1 (1-2)	0.86
Previous stillbirth	0 (0)	7 (3.1)	1.00
Previous neonatal death	1 (5)	8 (3.5)	0.54
Months since last pregnancy	24 (18-24)	20.5 (12-36)	0.83
Hypertensive disease of pregnancy	1 (5)	23 (10.1)	0.70
Severe pre-eclampsia	0 (0)	21 (9.2)	0.39
Eclampsia	2 (10)	8 (3.5)	0.19
Gestational diabetes	0 (0)	1 (0.4)	1.00
Number of fetus born	1 (1-1)	1 (1-1)	0.61
<i>Primary antenatal care provider</i>			0.003
<i>Dai</i>	2 (10.5)	1 (0.4)	
Auxiliary Nurse Midwife	1 (5.3)	2 (0.9)	
Primary Health Centre	1 (5.3)	17 (7.6)	
Community Health Centre	11 (57.9)	139 (61.8)	
Employees State Insurance facility	0 (0)	26 (11.6)	
Private practitioner	3 (15.8)	40 (17.8)	
Others	1 (5.3)	0 (0)	
Number of antenatal visits	4 (2-5)	5 (3-7)	0.05
Pregnancy at first ANC visit (mo)	3 (2-5)	3 (2-4)	0.62
Documented blood pressure during ANC	15 (75)	198 (87.2)	0.13
Documented hemoglobin during ANC	13 (65)	209 (92.1)	0.00
Documented urine examination during ANC	13 (65)	208 (91.6)	0.00
Documented fundal height	16 (80)	212 (93.4)	0.05
Took iron for 3 months	13 (65)	181 (79.7)	0.12
Distance between home and ANC site (km)	1 (0.5-2)	1 (0.5-2)	0.88

ANC: Antenatal care; Data represented as number (%), mean (SD) or median (interquartile range).