RESEARCH PAPER

Immunoglobulin Profile and Lymphocyte Subsets in Preterm Neonates

RAHUL SHARMA,¹ KANYA MUKHOPADHYAY,¹ AMIT RAWAT,¹ VANITA SURI,² SURJIT SINGH¹

From¹Departments of Pediatrics, and ²Obstetrics and Gynecology; Postgraduate Institute of Medical Education and Research, Chandigarh.

Correspondence to: Prof Kanya Mukhopadhyay, Division of Neonatology, Department of Pediatrics, Postgraduate Institute of Medical Education and Research, Chandigarh 160 012. kanyapgi@gmail.com Received: July 15, 2021; Initial review: September 21, 2021; Accepted: November 05, 2021. **Objective**: We documented the immunological profile of neonates and mothers, and lymphocyte subsets at birth. **Methods**: Consecutively born preterm neonates (26 to 31 weeks gestation) at our level III neonatal unit, fulfilling the inclusion criteria were enrolled. Immunoglobulin levels were assessed in maternal blood and in cord blood along with T cell subsets. **Results**: A total of 115 neonates were enrolled. The mean cord levels for IgG, IgM and IgA, respectively were 5.34, 0.10 and 0.04 g/L and of B, T, NK and NK-T cells were 14%, 71%, 10% and 1%, respectively of total lymphocyte population. Cord IgG and IgA levels showed a significantly rising trend with increasing gestation (P=0.005 and 0.02, respectively) but not IgM and T cell subsets. Maternal immunoglobulins were similar in all gestations. **Conclusion**: The cord IgG and IgA increased with increasing gestation but not IgM in neonates.

Keywords: B cells, Gestation, NK-T cells, T cells.

Published online: January 05, 2022; Pll: S097475591600389

Preterm neonates, born before 37 completed weeks of gestation, constitute 11% of total births with those born before 32 weeks comprising about 16% of preterm births [1,2]. Preterm neonates are seen to develop sepsis more often; one of the factors being immature immune system. Previous studies on immuno-globulin profile in preterm neonates have documented reduced immunoglobulin concentrations in preterm as compared to term babies. Studies on lymphocyte subsets and immunoglobulins in preterm neonates have had varying results [3-15]. Literature is scanty regarding immuno-globulin and lymphocyte subsets at very low gestation. Hence, our aim was to study the immunoglobulin profiles and lympho-cyte subsets in very preterm neonates at birth, which may have clinical implications.

METHODS

The study was conducted in a tertiary care referral hospital as a single center, observational study in neonates and their mothers after obtaining written informed consent from parents. The study was approved by the institute research ethics committee. All consecutively born neonates with gestation of 26 to 31 week during the study period of 1 year were enrolled. Neonates with severe perinatal asphyxia, suspected or proven chromosomal anomalies, intra-uterine infection and definite immunodeficiency in sibling or parent, and mothers with chorioamnionitis, multifetal gestation, acute febrile illness 4 weeks preceding delivery, on long-term steroids, recent vaccination and TORCH (toxoplasma, others, rubella, cytomegalovirus and herpes) infections were excluded. The gestational age of the baby was assessed from the first day of last menstrual period or first trimester ultrasonography, whichever was available. Postnatally gestational age was confirmed by New Ballard score. Demographic details, gravidity, mode of delivery, HIV (Human immunodeficiency virus) status, TORCH serology and any other infections during pregnancy were recorded for the mother. Neonatal details included birth weight, sex, APGAR and antenatal steroids. Various morbidities during the hospital stay including respiratory distress, sepsis, shock, total parenteral nutrition, neonatal jaundice, feed intolerance and hypoglycemia were recorded.

Each mother's venous blood (2 mL) was collected prior to delivery for immunoglobulin profile (IgG, IgM and IgA). Five mL of cord blood was collected for immunoglobulin profile (IgG, IgM and IgA) and lymphocyte subsets (B cells, T cells, NK cells and NK-T cells). Samples were centrifuged within 24 hours of collection and the separated serum was then stored in refrigerator at -80^oC and processed later for immunoglobulins. Lymphocyte subsets were assessed within 24 hours of collection. Serum IgG, IgA and IGM were estimated by endpoint nephelometry on a semi-automated nephelometer MININEPH (The Binding Site).

Lymphocyte subsets (T, B and NK, NK-T cells) were estimated using monoclonal antibodies against CD45, CD3, CD19 and CD16/56. Lymphocytes were gated using CD45 and side scatter, and the lymphocyte subsets were estimated in the gated lymphocyte population. Fifty μ L of EDTA anticoagulated blood was pipetted into a FACS tube and 10 μ L

INDIAN PEDIATRICS

each of fluorochrome labelled CD45, CD3, CD19 and CD56/ 16 were added to the tube and vortexed for proper mixing of the antibodies with the test sample. The tube was then incubated in the dark for 15 minutes. Following incubation, 1 mL of lysing solution was added to the tube and incubated in the dark for 10 minutes. The tube was then centrifuged at 1200 rpm for 10 minutes, the supernatant was discarded, and the stained cell pellets were washed twice with Phosphate buffer saline (PBS) and suspended in 500 µL of PBS before sample acquisition on a flow cytometer (Navios 2 laser 6 color flow cytometer, Beckman Coulter). Following sample acquisition, data analysis was performed using the Kaluza flow cytometry data analysis software. Lymphocytes were gated on CD45 vs side scatter and the different subsets in the gated lymphocytes were estimated using dot plots and histograms.

In the absence of previous data in very preterm neonates, it was decided to enrol a convenience sample of 25 mother-infant pairs at each gestational age.

Statistical analysis: Statistical analysis was done using SPSS version 22. Comparisons were made by using student *t* test, Mann-Whitney *U* test, Chi-square test and Kruskal-

Wallis test as appropriate. A P value of <0.05 was considered significant.

RESULTS

A total of 115 neonates and their mothers were enrolled in the study (26 to 27 weeks-18, 28 weeks-22 and 25 each from 29-31 weeks). The demographic details of the neonates are given in **Table I**. The immunoglobulin profile of neonates and their mothers are presented in **Web Tables I** and **Table II**. The lymphocyte subsets of the neonates are given in **Table III**.

The immunoglobulin profile (IgG, IgM and IgA) of mothers and neonates with and without sepsis did not differ significantly. The same was true for lymphocyte subsets in the neonates.

DISCUSSION

The present study noted increasing values of IgG and IgA with gestation. The lymphocyte subsets were not significantly different among the preterm neonates of 26 to 31 week gestation. There was no difference in immunoglobulin profile of neonates and mothers and lymphocyte subsets of

Table 1 Demographic and Chinical Characteristics of Freetin Neonates (N=115)							
Variables	26 wk (n=10)	27 wk (n=8)	28 wk (n=22)	29 wk (n=25)	30 wk (n=25)	31 wk (n=25)	
Birthweight (g) ^a	846 (169)	913 (121)	1076(198)	1112 (231)	1189 (217)	1369 (264)	
Male	5 (50)	5 (62)	13 (59)	10 (40)	13 (52)	11 (44)	
SGA	1 (10)	0	1 (5)	6(24)	9 (36)	8 (32)	
Suspect sepsis	7 (70)	5 (63)	12 (55)	18 (72)	20 (80)	10 (40)	
Proven sepsis	0	1(13)	2(9)	2(8)	1 (4)	1 (4)	

 Table I Demographic and Clinical Characteristics of Preterm Neonates (N=115)

Values in no. (%)or amean (SD). SGA-small for gestational age.

Variables (in g/L)	$26 wk (n{=}10)$	27 wk $(n=8)$	28 wk (n=22)	29 wk $(n=25)$	30 wk (n=25)	31 wk (n=25)
Immunoglobulin G	7.8 (6.2-11.9)	10.8 (9-12.8)	9.1 (6.8-11.2)	8.8 (7.2-9.9)	8.8 (6.4-10.9)	7.7 (7-9)
Immunoglobulin M	0.99 (0.5-1.4)	0.92 (0.8-1.1)	1.02 (0.8-1.6)	1.2 (0.9-1.5)	1 (0.5-1.5)	1.1 (0.8-1.3)
Immunoglobulin A	1.5 (1.1-1.8)	1.9 (1.6-2.4)	1.5 (1.2-2.0)	1.7 (1.2-2.1)	1.8 (1.3-2.3)	1.7 (1.2-1.9)

Values in median (IQR). All P > 0.05.

Table III Lymphocyte Subsets in	Cord Blood of Neonates With	1 Gestational Age of 26-31	Week (N=115
---------------------------------	-----------------------------	----------------------------	-------------

Variables	26 wk (n=10)	27 wk (n=8)	28 wk (n=22)	29 wk (n=25)	30 wk (n=25)	31 wk (n=25)
B cell	8.3 (5.1-13.7)	12.3 (8.9-19.2)	12.2 (8.3-22.2)	12.2 (9.4-19.3)	12.6 (8.5-19.1)	12.5 (8.7-18.1)
T cell	77.9 (73.6-84.3)	74.2 (64.7-78.7)	69.9 (61.6-79.5)	71.3 (58.1-80.1)	70.9 (59.9-79.8)	75.0 (65.6-80.8)
NK cell	7.2 (5.9-9.4)	8.4 (6.4-13.5)	10.2 (6.0-14.6)	9.0 (4.5-15.8)	10.4 (6.3-18.7)	7.4 (4.2-13.2)
NK-T cell	0.9 (0.5-1.7)	0.81 (0.5-2.1)	0.49 (0.4-1.0)	0.79 (0.5-1.2)	0.98 (0.4-1.3)	0.7 (0.6-1.1)

Values presented as percentage of total lymphocytes in median (IQR). All P>0.05. NK cell – natural killer cell.

WHAT THIS STUDY ADDS?

The normal values of immunoglobulins and lymphocyte subsets for neonates of each gestation are presented.

neonates who developed sepsis and who did not.

The immunoglobulin levels in the study were different from that by Boersma, et al. [4]. This was probably due to differences in the gestation ages of enrolled neonates, with lower gestation in present study compared to other studies. Ahmad, et al. [7] reported IgG levels in preterm in two groups (<34 weeks and >34 weeks). The IgG levels in the present study (5.34 g/L) are lower than Ahmad, et al. [7]. (6.41 g/L), the difference probably due to the differences in gestation in enrolled neonates. The study by Ozdemir, et al. [6] enrolled neonates of <28 weeks, 29-31 weeks, 32-37 weeks and ≥38 weeks. The mean (min-max) IgG levels were 3.7 (1.5-9.6) g/L, 5.4 (3.1-8.7) g/L, 6.7 (3.3-11) g/L and 7.9 (4-20) g/L, respectively. Conway, et al. [13] noted a linear correlation of IgG levels with increasing gestation. Our study also showed similar trend. Sharma, et al. [14] found mean (SD) and range of IgG, IgM and IgA values in 40 preterm infants were 11 (1.5) g/L and 9-14 g/L, 0.34 (0.06) g/L and 0.2 to 0.4 g/L, 0.02 (0.03) g/L and 0-0.12 g/L, respectively, which are higher than our study, probably because they included preterm infants of higher gestations up to 36 weeks. Panayotou, et al. [15] studied serial IgA and IgM in cord blood in healthy preterm neonates of 28-35 weeks of gestation and found that IgA was absent in most cord blood samples. This was in contrast to the presence of IgA in cord blood samples in the present study. As IgM does not cross placenta and hence high cord IgM may reflect underlying intrauterine infection however we had very low value of IgM in our study.

When comparing lymphocyte subsets, there were no studies that had separate values at each gestation. Berrington, et al. [8], observed that preterm had 65% T cells, 23% B cells and 7% NK cells in their blood. These were nearly similar to our study. The T cells and B cells in the study by Quinello, et al. [9] in gestation 30 weeks to 33 weeks were 43.3% and 16.1%, respectively. Our study showed T cells and B cells to be 70.8% and 13.9%, respectively at gestation 30 to 31 weeks. Our study also found no difference in lymphocyte subsets in those who developed sepsis and those who did not. This is in contrast to the study by Bochennek, et al. [12], which showed that neonates who developed sepsis had significantly lower NK cells. The cause for this could not be evaluated.

This study showed that there was no statistically significant difference in the immunoglobulin profile and

lymphocyte subsets of the preterm neonates with regards to occurrence of sepsis. However, this study had a small sample size and further studies are needed to establish the normal values of immunoglobulins and lymphocyte subsets at each gestation and their relation to occurrence of sepsis.

Ethics clearance: Approved by the institutional ethics committee no. NK/3996/MD; dated June 28, 2019.

Contributors: KM, RS: conceived the idea of this study; RS, KM, AR, VS, SS: involved in formulating the protocol of this study; RS: collected the samples for this study. The samples were processed in the laboratory under guidance of AR and SS. RS, KM, AR, VS, SS were involved in the writing of this manuscript. All authors approved the final version of manuscript, and are accountable for all aspects related to the study.

Funding: None; Competing interests: None stated.

Note: Additional material related to this study is available with the online version at *www.indianpediatrics.net*

REFERENCES

- Melville JM, Moss TJM. The immune consequences of preterm birth. Front Neurosci. 2013;7:1-9.
- Blencowe H, Cousens S, Chou D, et al. Born too soon: The global epidemiology of 15 million preterm births. Reprod Health. 2013;10:S2
- Berg T. Immunoglobulin levels in infants with low birth weights. Acta Paediatrica. 1968;57:369-76.
- 4. Boersma ER. Serum immunoglobulins IgG, IgM, and IgA in maternal cord blood pairs from infants of normal and low birthweights in Tanzania. Arch Dis Child. 1981;56:31-5.
- Sierig G, Labitzke B, Diez U, et al. Natural history of serum immunoglobulin concentrations in low birth weight infants and association with respiratory tract infections. Biol Neonate. 2002;82:159-65.
- Alkan Ozdemir S, Ozer EA, Kose S, et al. Reference values of serum IgG and IgM levels in preterm and term newborns. J Matern Fetal Neonatal Med. 2016;29:972-76.
- Ahmad P, Hameed F, Ahmad KN. Serum IgG levels in full term preterm and SFD neonates. Indian J Pediatr. 1986; 53:405-7.
- Berrington JE, Barge D, Fenton AC, et al. Lymphocyte subsets in term and significantly preterm UK infants in the first year of life analysed by single platform flow cyto-metry. Clin Exp Immunol. 2005;140:289-92.
- Quinello C, Silveira-Lessa AL, Ceccon MEJR, et al. Phenotypic differences in leucocyte populations among healthy preterm and full-term newborns. Scand J Immunol. 2014;80:57-70.
- 10. Mukhopadhyay D, Weaver L, Tobin R, et al. Intrauterine growth restriction and prematurity influence regulatory T cell

INDIAN PEDIATRICS

development in newborns. J Pediatr Surg. 2014;49: 727-32.

- 11. Kent A, Scorrer T, Pollard AJ, et al. Lymphocyte subpopulations in premature infants: an observational study. Arch Dis Child Fetal Neonatal Ed. 2016;101:F546-51.
- 12. Bochennek K, Fryns E, Wittekindt B, et al. Immune cell subsets at birth may help to predict risk of late-onset sepsis and necrotizing enterocolitis in preterm infants. Early Hum Dev. 2016;93:9-16.
- 13. Conway SP, Dear PR, Smith I. Immunoglobulin profile of the preterm baby. Arch Dis Child. 1985;60:208-12.
- 14. Sharma S, Lal H, Saigal RK. Immunoglobulins IgG, IgM and IgA levels in preterm and small for date newborns. Indian Pediatr.1991;28:741-44
- 15. Panayotou P, Papadatos C, Papaevangelou G, et al. Immunoglobulin A and M levels in premature infants with gastroenteritis. Arch Dis Child. 1971;46:671-75.

Variables	26 wk	27 wk	28 wk	29 wk	30 wk	31 wk
	n=10	n=08	n=22	n=25	n=25	n=25
Immunoglobulin (g/L) ^a	4	3.7	4.7	5.1	5.6	5.3
	(2.7-4.8)	(3.3-6.7)	(4-5.6)	(4.2-6.3)	(4.8-7.2)	(4.6-7.2)
Immunoglobulin (g/L) ^b	0.09	0.07	0.05	0.06	0.06	0.09
	(0.04-0.2)	(0.04-0.08)	(0.04-0.1)	(0.04-0.08)	(0.04-0.08)	(0.07-0.1)
Immunoglobulin						
(g/L) ^a	0.03	0.03	0.03	0.03	0.03	0.04
	(0.03-0.05)	(0.03-0.03)	(0.03-0.03)	(0.03-0.05)	(0.03-0.03)	(0.03-0.06)

Web Table I Immunoglobulin Levels in Cord Blood of Neonates With Gestational Age of 26-31 Week (N=115)

Values in median (IQR). ${}^{a}P < 0.05$; ${}^{b}P = 0.051$.