

report recommends serial repetition of TFT in ELBW neonates, particularly those having initial stormy course during hospitalisation.

**SANTOSH KUMAR PANDA, SUSHREE SMITA BEHURA,  
DEEPTI DAMAYANTY PRADHAN\***

*Department of Paediatrics, KIMS,  
Bhubaneswar, Odisha  
deepti.pradhan@kims.ac.in*

## REFERENCES


1. Hashempour M, Hovsepian S, Ansari A, et al. Screening of congenital hypothyroidism in preterm, low birth weight and very low birth weight neonates: A systematic review. *Pediatr Neonatol.* 2018;59:3-14.
2. Sudhanshu S, Riaz I, Sharma R, et al. Newborn screening guidelines for congenital hypothyroidism in India: Recommendations of the Indian Society for Pediatric and Adolescent Endocrinology (ISPAE) - Part II: Imaging, treatment and follow-up. *Indian J Pediatr.* 2018;85:448-53.
3. Léger J, Olivieri A, Donaldson M, et al. Congenital Hypothyroidism Consensus Conference Group. European Society for Paediatric Endocrinology Consensus Guidelines on Screening, Diagnosis, and Management of Congenital Hypothyroidism. *J Clin Endocrinol Metab.* 2014;99:363-84.
4. Mandel SJ, Hermos RJ, Larson CA, et al. Atypical hypothyroidism and the very low birthweight infant. *Thyroid.* 2000;10:693-5.
5. Kaplowitz PB. Neonatal thyroid disease: testing and management. *Pediatr Clin North Am.* 2019;66:343-52.
6. Zung A, Bier Palmon R, Golan A, et al. Risk factors for the development of delayed TSH elevation in neonatal intensive care unit newborns. *J Clin Endocrinol Metab.* 2017;102: 3050-55.

## Adenomyomatosis of Gallbladder in a Neonate

Adenomyomatosis of gallbladder is defined as hypertrophy of gallbladder mucosal epithelium that invaginate into thickened muscularis propria, leading to formation of intramural diverticula. It is mainly seen in adults and the incidences increases with increasing age, as most of the documented cases are above 50 years of age, and many were detected in cholecystectomy specimens. Adenomyomatosis of gallbladder is very rare in the paediatric population. We herein report a case of adenomyomatosis of gallbladder in a newborn infant.

A term infant was delivered to a primigravida mother at 37 weeks of gestation at our hospital. Post-natal transition of the baby was smooth. We noted that one of the antenatal ultrasonographic scan of the mother was suggestive of umbilical varices in the fetus. To detect any associated abnormality in the neonate due to the umbilical arices, a postnatal ultrasound was done. The scan showed a partially distended gallbladder with diffuse wall thickening and multiple tiny echogenic foci in anterior and posterior walls involving entire gallbladder showing comet tail artefacts (**Fig.1**). The imaging findings were consistent with the adenomyomatosis of gallbladder. The baby continued to do well clinically and discharged with the advice to repeat the abdominal USG later.

The exact pathophysiology of adenomyomatosis of gallbladder is not clearly known. One theory suggests that change in intra-cystic pressure because of disruption of gallbladder function may lead to proliferation of cells in the gallbladder mucosa and hyperplasia of the muscle layer. The epithelial layer can then invaginate into the muscular layer, leading to the formation of Rokitsky Ashoff Sinuses (RAS) [1]. This is also seen on histopathology, which generally shows hyperplasia of the epithelium and mucosal out pouching through the muscular layer [2].



**Fig. 1** Ultrasonography of abdomen showing adenomyomatosis of gall bladder.

Most of the cases in adults were diagnosed when they developed cholelithiasis or any other gall bladder condition. Although, there are some reports suggesting an association between adenomyomatosis and gallbladder cancer, there is no concrete evidence to prove this association [3].

A review of the literature showed only nine reported cases of paediatric gallbladder adenomyomatosis, with ages ranging from 1 day to 14 years. The most common symptom was abdominal pain, which was seen in 7 of the patients. Cholecystectomy was performed in seven paediatric cases [4]. One patient was a 1-day old newborn baby with suspected heterotaxy syndrome and complete atrioventricular canal. Patients generally did not have other significant comorbid conditions [5].

In children, adenomyomatosis of gall bladder remains an incidental finding on USG scan for some other reason as there are no specific clinical symptoms. Computed tomography and magnetic resonance imaging have also been used in the diagnosis of this

condition [1]. Usually, no treatment is recommended. cholecystectomy is required in patient who are symptomatic with right upper quadrant pain.

**JP DADHICH,<sup>1\*</sup> P KUMAR,<sup>1</sup> S MITTAL,<sup>2</sup> CP DADHICH<sup>3</sup>**

<sup>1</sup>*Paediatrics and Neonatology, <sup>2</sup>Radio-diagnosis,*

<sup>3</sup>*Obstetrics and Gynaecology,*

*Fortis Escorts Hospital, Jaipur, Rajasthan.*

*jpdadhich@gmail.com*

## REFERENCES

1. Golse N, Lewin M, Rode A, et al. Gallbladder adenomyomatosis: Diagnosis and management. *J Visc Surg.* 2017;154:

345-53.

2. Akçam M, Buyukyavuz I, Ciris M, et al. Adenomyomatosis of the gallbladder resembling honeycomb in a child. *Eur J Pediatr.* 2008;167:1079-81.
3. Pang L, Zhang Y, Wang Y, et al. Pathogenesis of gallbladder adenomyomatosis and its relationship with early-stage gallbladder carcinoma: An overview. *Braz J Med Biol Res.* 2018;51:e7411.
4. Alapati S, Braswell LE. Neonatal adenomyomatosis of the gallbladder: An incidental finding at 12 hours of life. *Radiol Case Rep.* 2014;9:859.
5. Agrusti A, Gregori M, Salviato T, et al. Adenomyomatosis of the gallbladder as a cause of recurrent abdominal pain. *J Pediatr.* 2018;202:328.

## Thiamine-Responsive Acute Pulmonary Hypertension in an Exclusively Breastfed Infant

Thiamine deficiency presenting as acute severe pulmonary hypertension (PAH) in exclusively breastfed babies has been reported from various parts of India recently [1-3]. Here we report a male infant who presented with acute pulmonary hypertension and responded to thiamine.

A 4-month-old male infant, first born to non-consanguineous parents, belonging to the Chetty community from Mysuru district of Karnataka was referred to our hospital in view of acute onset of cough and respiratory distress. There was a history of approximately ten recent deaths in their community in babies of the same age group, one of them being this baby's first cousin. Most of them presented with bronchiolitis like symptoms, rapidly progressing to acute respiratory distress syndrome (ARDS). This baby was referred to our hospital with the possibility of inborn errors of metabolism.

Baby was delivered at term, was on exclusive breast feeds, thriving well (birth weight 3 kg, present weight 6.5 kg) and had normal development. At presentation, the baby had tachypnea, tachycardia, hypoxia (SpO<sub>2</sub> 92% in room air) and respiratory distress. Liver was palpable 3 cm below costal margin. Second heart sound was loud and wide split. There was hypotonia and deep tendon reflexes were absent. The arterial blood gas showed mixed disorder with metabolic acidosis and respiratory alkalosis. Tandem mass spectroscopy was normal. Electrocardiogram showed evidence of right atrial (RA) dilatation and right ventricular (RV) hypertrophy (RVH). Echocardiography showed features of severe PAH (RA, RV dilated, moderate tricuspid regurgitation, inter-atrial septum bulging into left atrium) without any evidence of structural heart disease. Contrast enhanced computed tomography (CECT) of lung showed pulmonary plethora with features of PAH without any intrinsic pulmonary pathology.

A detailed dietary history revealed that mother's diet mainly consisted of polished white rice and *Rasam* (a sour soup made

with tomato and lentil). There was traditional avoidance of other foods like pulses, vegetables, and non-vegetarian foods during postpartum period. Considering the probability of thiamine deficiency, thiamine was started in a dose of 10 mg/kg, orally. The baby showed clinical improvement, could be weaned off oxygen by 12 hours, and respiratory distress subsided by 48 hours. A repeat echocardiography at 48 hours showed marked improvement with only mild PAH, and subsequently showed complete resolution of PAH at 96 hours. No pulmonary vasodilators other than oxygen were given.

The whole blood thiamine level estimated by HPLC of both baby and mother were low (25 µg/mL and 27.25 µg/mL, respectively, normal range being 28 to 85 µg/mL). There was no history of aphonia, encephalopathy, seizures, or oedema. Serum 25-hydroxy vitamin D3 level was normal. The history, geographic background, clinical features, investigations, and dramatic response to thiamine confirmed the diagnosis of Thiamine responsive acute pulmonary hypertension of early infancy (TRAPHEI). Family was counselled about the dietary modification and both mother and baby were put on thiamine supplementation at 2 mg/kg/day. A repeat echocardiography at one month follow-up visit showed complete recovery. ECG normalized by two months. Muscle tone and deep tendon reflexes normalised after two months of thiamine therapy, suggestive of coexistent neuropathic form of thiamine deficiency. Neuroimaging was not done, as the baby had improved.

Thiamine has an important role in the metabolism of carbohydrates and synthesis of nucleic acid, myelin, and neurotransmitters. A regular dietary supply is must for preventing deficiency because of high turnover rate and low body stores of thiamine. Thiamine causes preferential injury to tissues with heavy metabolic turnover like neurons and cardiac myocytes. The clinical spectrum of thiamine deficiency in children can be acute cardiac form, aphonic form, pseudo meningitic form or neuropathic form [4]. Thiamine deficiency has been documented in those consuming polished cereals and with other dietary restrictions. Infants are at highest risk due to rapid growth rate and high energy needs. In exclusively breastfed babies of thiamine deficient mothers, thiamine levels rapidly decline in the third month of life. As in our case, culturally determined food restrictions during the postpartum period are prevalent in various parts of India [5].